

# Boundedness and regularity of the solutions of a kind of nonlinear systems

Lubomira Softova

University of Salerno, Italy

E-mail: lsoftova@unisa.it

We deal with nonlinear systems of elliptic and parabolic type

$$\operatorname{div} \mathbf{A}(x, \mathbf{u}, D\mathbf{u}) + \mathbf{b}(x, \mathbf{u}, D\mathbf{u}) = 0 \quad x \in \Omega$$

$$\mathbf{u}_t - \operatorname{div} \mathbf{A}(x, t, \mathbf{u}, D\mathbf{u}) + \mathbf{b}(x, t, \mathbf{u}, D\mathbf{u}) = 0 \quad (x, t) \in Q_T = \Omega \times (0, T)$$

satisfying componentwise structural conditions. The nonlinear terms are Carathéodory maps having controlled growth with respect to the solution and the gradient and the data are in suitable Lebesgue and Morrey spaces.

Under these assumptions we obtain essential boundedness of the weak solutions and in the elliptic case also optimal Morrey regularity for the solutions.

## REFERENCES

- [1] E.A. ALFANO, L. FATTORUSSO, L.G. SOFTOVA, Boundedness of the solutions of a kind of non-linear parabolic systems, *J. Differ. Equ.* **360** (2023), 51–66.
- [2] L. FATTORUSSO, L. SOFTOVA, Precise Morrey regularity of the weak solutions to a kind of quasi-linear systems with discontinuous data, *Electron. J. Qual. Theory Diff. Equ.*, **36** (2020) 13p.
- [3] D.K. PALAGACHEV, L. SOFTOVA, Boundedness of solutions to a class of coercive systems with Morrey data, *Nonlin. Anal.* **191** (2020), 111630.
- [4] L. SOFTOVA, Boundedness of the solutions to non-linear systems with Morrey data, *Compl. Var. Elliptic Equ.* **63**, no. 11 (2018), 1581–1594.